DRAFT

Fire Regime Condition Class (FRCC) Interagency Handbook Reference Conditions

Modeler: KellyAnn Gorman Date: 20 October PNVG Code: NHDW1

2004

Potential Natural Vegetation Group: Northern Hardwood Forest

Geographic Area: Northern tier of states in the east, including New England, NY, and the northern parts of PA, MI, and WI; also a southern extension in the Appalachian Mountains, including NC, TN, VA, and WV, and south to northern GA.

Description: Tall, broadleaf deciduous forest with few evergreen trees occurring on high-elevation, concave landforms and north-facing slopes in high mountain areas. Dominated by sugar maple (*Acer saccharum*), beech (*Fagus grandifolia*), yellow birch (*Betula allegheniensis*), northern red oak (*Quercus rubra*), and black cherry (*Prunus serotina*). Other common associates include buckeye (*Aesculus flava*), striped maple (*A. pensylvanicum*), red maple (*A. rubrum*), mountain maple (*A. spicatum*), white ash (*Fraxinus americana*), hop hornbeam (*Ostrya virginiana*), mountain-ash (*Sorbus americana*), basswood (*Tilia americana*), serviceberry (*Amelanchier arborea*), dogwood (*Cornus alternifolia*), and American elm (*Ulmus americana*).

Fire Regime Description: Fire Regime Group V. Fire disturbances are severe and affect large patch sizes but are very rare, at 300 to 1,000-year intervals; wind events are much more frequent. Other disturbances, including windthrow, insect attack, and ice storms, usually on a single-treegap scale, are more important than fire although they may have pre-disposed the forest to fire during drought conditions.

Vegetation Type and Structure

Class*	Percent of	Description
	Landscape	
A: post replacement	10	Young stand characterized by yellow birch, red maple, ash, and mountain-ash; less than 25 yrs old
B : mid-seral closed	30	Intermediate stand characterized by northern red oak, black cherry, red maple, and white ash; 25 - 100 yrs old
E: late- seral closed	60	Mature stand dominated by sugar maple, beech, and yellow birch; over 100 yrs old
Total	100	•

^{*}Formal codes for classes A-E are: AESP, BMSC, CMSO, DLSO, and ELSC, respectively.

Fire Frequency and Severity

_ increquency and coroning							
	Fire Frequency	Probability	Percent,	Description			
Fire Severity	(yrs)	-	All Fires				
Replacement Fire	650	.0015	94				
Non-Replacement Fire	10,000	.0001	6				
All Fire Frequency*	650	.0016	100				

^{*}All Fire Probability = sum of replacement fire and non-replacement fire probabilities. All Fire Frequency = inverse of all fire probability (previous calculation).

References

Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.

Kuchler, A.W. 1964. Northern hardwoods (*Acer-Betula-Fagus-Tsuga*). #106 *In:* Manual to accompany the map Potential Natural Vegetation of the United States. New York, NY: The American Geographical Society. 156 p.

Loehle, Craig. 1988. Tree life histories: the roles of defenses. Canadian Journal of Forest Research 18: 209-222.

Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.

Southern Appalachian Man and the Biosphere. 1996. The Southern Appalachian Assessment Terrestrial Technical Report. Report 5 of 5. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region.

US Department of Agriculture, Forest Service, Southern Region. 1997. Montane and allied spruce and spruce-fir forest old-growth forest community. Pp. 100-102 *in*: Guidance for conserving and restoring old-growth forest communities in National Forests in the Southern Region: Report of the Region 8 Old-Growth Team. Forestry Report R8-FR 62. Atlanta, GA: U.S. Department of Agriculture, Forest Service, Southern Region. 120 p.

U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online 12 February 2004]. Available: http://www.fs.fed.us/database/feis/.

PERSONAL COMMUNICATION (if applicable):

VDDT File Documentation

Assumptions: Patch-size scale is single-tree canopy gaps.

Native American fire was considered but was not determined to be a significant factor.













